Critical Behavior of a Relativistic Bose Gas

P. N. Pandita
P - 30, North Eastern Hill University, Shillong 793 022, India

We show that the thermodynamic behavior of relativistic ideal Bose gas, recently studied numerically by Grether et al., can be obtained analytically. Using the analytical results, we obtain the critical behavior of the relativistic Bose gas exactly for all the regimes. We show that these analytical results reduce to those of Grether et al. in different regimes of the Bose gas.

PACS numbers: 03.75.Kk, 05.30.Jp, 05.30.-d

arXiv:1406.7389v1 [cond-mat.stat-mech] 28 Jun 2014

Bose-Einstein condensation (BEC) of an ideal Bose gas has been a subject of extensive studies. In particular Bose-Einstein condensation in a relativistic ideal Bose gas with nonzero chemical potential has been studied by several authors [1–5]. In earlier papers [1, 6] on relativistic ideal Bose condensation, antiboson production was not taken into account. The necessity of the antiboson contribution to the thermodyanmics of Bose gas at relativistic energies was pointed out in [3], and a high-temperature expansion for various thermodynamic quantities was established and studied. At sufficiently high temperatures, antibosons are expected to be pair produced in sufficient numbers so that their contribution cannot be neglected.

In a recent paper, Grether et al. studied Bose-Einstein condensation in a relativistic ideal Bose gas, and calculated its thermodyanamic properties numerically [7]. In this note we point out that the model can be solved exactly in a closed form, including antibosons, without making any approximations, and show that an analytic expression for thermodynamic quantities can be obtained at all temperatures and the critical behavior studied in general. The analytical study is physically more transparent, and sheds light on the connection between relativistic and the non-relativistic ideal Bose gas by showing that the critical behavior is same for the two systems. It is only the overall amplitude which is different for the two systems. We also show that our exact results reduce to the results of ref. [7] in different regions studied by these authors.

The ideal Bose gas is characterized by three basic length scales, namely the thermal wavelength λ_T, the mean interparticle spacing $\overline{\lambda}$, and the Compton wavelength λ_C. From these, one can obtain two independent ratios which we consider to be $R_1 = \overline{\lambda}/\lambda_T$, and $R_2 = \lambda_C/\lambda_T$. For quantum (classical) gas we have $R_1 \ll 1$ ($\gg 1$), whereas for $R_2 \ll 1$ and $R_2 \gg 1$ we will have nonrelativistic or ultrarelativistic gas, respectively. Clearly, it is important to understand the connection between different regions, characterized by different length scales, and have a unified treatment for all the regions involved. In this note we solve the model exactly in general, which includes all the cases corresponding to all the length scales and without making a high temperature expansion. We show analytically that the critical behavior of the model is the same as that of standard nonrelativistic Bose gas. The difference between nonrelativistic and the ultrarelativistic gas, where antiparticles are important, arises only in the values of critical amplitudes.

The net "charge" or number density, assumed positive without loss of generality, for a relativistic Bose gas of N bosons and $\overline{N}$ antibosons, each of mass m, can be written as (the notation is standard, and we choose units $\hbar = c = k_B = 1, \beta = T^{-1}$; g is the spin degeneracy factor)

$$n = g \int \frac{d^3k}{(2\pi)^3} \left[\frac{1}{\exp[\beta(k^2+m^2)^{1/2} - \beta\mu] - 1} - \frac{1}{\exp[\beta(k^2+m^2)^{1/2} + \beta\mu] - 1} \right], \tag{1}$$

where we have enclosed the system in a cubical box of volume L^3, with L being the length of an edge in each of the 3 spatial dimensions. This can be further written as

$$n = \frac{m^3 g}{2\pi^2} \int_0^\infty dx \;\; x^2 \;\; \frac{\sinh(\alpha - \phi)}{\cosh a(x) - \cosh(\alpha - \phi)} \equiv \frac{m^3 g}{2\pi^2} \mathcal{W}_3(\alpha, \phi), \tag{2}$$

where $\alpha = \beta m$, $a(x) = \alpha(1 + x^2)^{1/2}$, and $\phi = -\beta(\mu - m)$. The parameter ϕ is defined so as to reduce to its nonrelativistic counterpart in that limit. For d spatial dimensions this generalizes to

$$n = a_d^{-1} \mathcal{W}_d(\alpha, \phi), \tag{3}$$

where

$$a_d = 2^{d-1} \pi^{d/2} \Gamma(d/2) m^{-d} g^{-1}, \tag{4}$$

$$\mathcal{W}_d(\alpha, \phi) = \int_0^\infty dx \;\; x^{d-1} \;\; \frac{\sinh(\alpha - \phi)}{\cosh a(x) - \cosh(\alpha - \phi)}. \tag{5}$$

The results (3), (4) and (5) agree with the result (8) of ref.[7]. Similarly, we can write the energy density as

$$u \;=\; g\int \frac{d^3k}{(2\pi)^3}\left[\frac{(k^2+m^2)^{1/2}}{\exp[\beta(k^2+m^2)^{1/2}-\beta\mu]-1}+\frac{(k^2+m^2)^{1/2}}{\exp[\beta(k^2+m^2)^{1/2}+\beta\mu]-1}\right], \tag{6}$$

which can be rewritten as

$$u \;=\; ma_d^{-1}\mathcal{Y}_d(\alpha,\phi), \tag{7}$$

where

$$\mathcal{Y}_d(\alpha,\phi) \;=\; \int_0^\infty dx \;\; x^{d-1} \;\; \frac{\alpha^{-1}a(x)\cosh(\alpha-\phi)-\sinh(\alpha-\phi)-\alpha^{-1}a(x)e^{-a(x)}}{\cosh a(x)-\cosh(\alpha-\phi)}. \tag{8}$$

In order to study the critical behaviour of relativistic Bose gas, it is useful to perform the integrals in (5) and (8) in a closed form by expanding the integrands in infinite series and introducing modified Bessel functions in the resulting integrals. Performing this, we get

$$\mathcal{W}_d(\alpha,\phi) \;=\; b_d\sum_{r=1}^{\infty} r^{-(d'-1)}\sinh(r\alpha-r\phi)K_{d'}(r\alpha), \tag{9}$$

$$\mathcal{Y}_d(\alpha,\phi) \;=\; b_d\sum_{r=1}^{\infty} r^{-(d'-1)}[\cosh(r\alpha-r\phi)K_{d'+1}(r\alpha)-\sinh(r\alpha-r\phi)K_{d'}(r\alpha)-(r\alpha)^{-1}\cosh(r\alpha-r\phi)K_{d'}(r\alpha)] \tag{10}$$

where $K_{d'}(r\alpha)$ is a modified Bessel function of order d' and real argument $r\alpha$ [8, 9], and

$$b_d \;=\; \pi^{-1/2}2^{d'}\Gamma(d/2)\overline{\alpha}^{(d'-1)}, \qquad\qquad d'=\frac{1}{2}(d+1). \tag{11}$$

We can study the critical behavior of the relativistic Bose gas by studying the variation of μ, or equivalently, ϕ as a function of T. By standard argument of keeping charge density positive, μ must satisfy the condition $-m\le\mu\le m$. The parameter ϕ is, therefore, defined to be positive. From Eqs.(5) or (9) it is clear that as $\beta\to 0, \mu\to 0$, or $\phi\to\beta m$. Decreasing β, increases μ until it reaches its limiting value m. The critical temperature is, thus, obtained from (3) for $\phi=0$. Thus,

$$a_d n \;=\; \mathcal{W}_d(\alpha_c,0)=b_{dc}\sum_{r=1}^{\infty} r^{-(d'-1)}\sinh(r\alpha_c)K_{d'}(r\alpha_c), \tag{12}$$

where b_{dc} is the value of b_d at $\alpha_c=\beta_c m=m/T_c$. Using the large argument form of the function $K_\nu(z)$

$$K_\nu(z) \;\approx\; [\frac{\pi}{2z}]^{1/2}e^{-z}[1+\frac{4\nu^2-1}{8z}+.....], \tag{13}$$

we see that the ratio of the successive terms of the series (12) can be written as

$$\frac{u_{r+1}}{u_r} \;=\; 1-\frac{d}{2r}+\mathcal{O}(\frac{1}{r^2}), \qquad r\to\infty, \tag{14}$$

so that by ratio test the series converges for $d>2$. We, therefore, conclude that a nonzero T_c exists. *It is important to note that this result is independent of* $\alpha_c(\equiv mc^2/(k_BT_c))$.

Using (3) and (5) we can determine the behavior of ϕ in the critical region. To do so, we expand $\mathcal{W}_d(\alpha,\phi)$ near $\phi=0$, which can be obtained by calculating $\partial\mathcal{W}_d(\alpha,\phi)/\partial\phi$ at $\phi=0$. We obtain

$$\partial\mathcal{W}_d(\alpha,\phi)/\partial\phi|_{\phi=0} \;=\; -b_d\sum_{r=1}^{\infty} r^{-(d'-2)}\cosh(r\alpha)K'_d(r\alpha). \tag{15}$$

This sum converges only for $d>4$. The behavior for $2<d<4$ as $\phi\to 0$ can be obtained by calculating the derivative

$$\partial\mathcal{W}_d(\alpha,\phi)/\partial\phi \;=\; -b_d\sum_{r=1}^{\infty} r^{-(d'-2)}\cosh(r\alpha-r\phi))K'_d(r\alpha), \tag{16}$$

from which we get, asymptotically,

$$\partial \mathcal{W}_d(\alpha,\phi)/\partial\phi \;\approx\; -\frac{1}{2}(2/\alpha)^{d/2}\Gamma(d/2)F_{(d-2)/2}(\phi), \tag{17}$$

$$F_n(\phi) \;=\; \sum_{r=1}^{\infty} r^{-n}e^{-r\phi}, \tag{18}$$

$$F_n(\phi) \;\approx\; \Gamma(1-n)\phi^{n-1}, \qquad n<1, \phi\to 0. \tag{19}$$

$$\tag{20}$$

Putting these results together, we can write

$$\partial \mathcal{W}_d(\alpha,\phi)/\partial|\phi| \;\approx\; -\frac{1}{2}(2/\alpha)^{d/2}\Gamma(d/2)\Gamma(2-d/2)\phi^{(d-4)/2}, \tag{21}$$

for $2<d<4$. However, the quantity $\partial\mathcal{W}_d(\alpha,0)/\partial\alpha$ is calculated to be finite and negative at α_c. We can , therefore, write

$$\mathcal{W}_d(\alpha,0) \;\approx\; \mathcal{W}(\alpha_c,0)-(\alpha-\alpha_c)\mathcal{W}', \tag{22}$$

$$\mathcal{W}' \;\equiv\; -[d\mathcal{W}_d(\alpha,0)/d\alpha]_{\alpha=\alpha_c}. \tag{23}$$

Using (21) and (22) in (3), we can write

$$\phi \;\approx\; (C^+)^{-1}t^{2/(d-2)}, \qquad 2<d<4, \qquad t>0, \tag{24}$$

$$C^+ \;=\; [\frac{2^{d/2}\Gamma(d/2)\Gamma(2-d/2)}{(d-2)\alpha_c^{(d+2)/2}\mathcal{W}'}]^{2/(d-2)}, \tag{25}$$

where we have written

$$t \;=\; (T-T_c)/T_c. \tag{26}$$

From the result (24) we note that the critical exponent is $2/(2-d)$, which is the same as that of the nonrelativistic Bose gas. Thus, the critical behavior of the gas is same in all regions, whether relativistic or nonrelativistic [10]. The only difference relates to the amplitude, the overall factor multiplying $[(T-T_c)/T_c]^{2/(d-2)}$ in (24). Finally, turning to the behavior for $T<T_c$, we note that $\phi=0$ for $T<T_c$.

In order to see the connection of our general analytical results with the results of ref. [7], we consider the nonrelativistic (NR) and the ultrarelativistic (UR) limit for the number density and the energy density of the Bose gas. These correspond to $\alpha_c \gg 1$ and $\alpha_c \ll 1$, respectively. From Eq.(12), we have

$$\mathcal{W}_d(\alpha_c,0) \;=\; b_{dc}\sum_{r=1}^{\infty} r^{-(d'-1)}\sinh(r\alpha_c)K_{d'}(r\alpha_c). \tag{27}$$

Using (13), we get for the nonrelativistic case

$$\mathcal{W}_d(\alpha_c,0) \;=\; \frac{1}{2}\left[\frac{2}{\alpha_c}\right]^{d/2}\Gamma(d/2)\zeta(d/2), \qquad \alpha_c\gg 1. \tag{28}$$

Using the fact that $\mathcal{W}_d(\alpha_c,0)=a_d n$, and solving (28) for the critical temperature $\alpha_c=m/T_c$, we get

$$T_c \;=\; \frac{2\pi}{m}\left[\frac{n}{g\zeta(d/2)}\right]^{2/d}, \tag{29}$$

in the nonrelativistic region, which is the result (5) of ref.[7]. Using (7), (8), and (10), we can write for the energy density in the nonrelativistic region

$$u \;=\; ma_d^{-1}\mathcal{Y}_d(\alpha_c,0), \tag{30}$$

where

$$\mathcal{Y}_d(\alpha_c,0) \;=\; \frac{1}{2\alpha_c}\left[\frac{2}{\alpha_c}\right]^{d/2}\Gamma(\frac{d+2}{2})\zeta(d/2), \qquad \alpha_c\gg 1. \tag{31}$$

On the other hand for the ultrarelativistic limit $\alpha_c \ll 1$, we can use the expansion [9]

$$K_\nu(z) \approx \frac{2^{\nu-1}\Gamma(\nu)}{z^\nu}, \tag{32}$$

for $z \ll 1$ to obtain

$$\mathcal{W}_d(\alpha_c, 0) = \frac{2}{\alpha_c^{d-1}}\Gamma(d)\zeta(d-1), \qquad \alpha_c \ll 1 \tag{33}$$

which, together with the result (12), can be solved for the critical temperature to obtain

$$T_c = \left[\frac{n(2\pi)^d\Gamma(d/2)}{4gm\pi^{d/2}\Gamma(d)\zeta(d-1)}\right]^{1/(d-1)}, \qquad \alpha_c \ll 1, \tag{34}$$

which is the same as the result (9) of ref. [7]. Finally, in the UR limit the energy density can be written as

$$u = ma_d^{-1}\mathcal{Y}_d(\alpha_c, 0), \tag{35}$$

$$\mathcal{Y}_d(\alpha_c, 0) = \frac{2}{\alpha_c}(d-1)\Gamma(d)\zeta(d-1), \qquad \alpha_c \ll 1. \tag{36}$$

In summary we have solved the ideal, relativistic Bose gas, in a closed form, and have shown that the critical behavior of the gas is same as for the standard nonrelativistic Bose gas. Our general results are applicable to all the regions of the Bose gas. We have shown that our general results reduce to the known results in the nonrelativistic and the ultrarelativistic regions. We have, thus, obtained a unified treatment of the Bose gas which is applicable in all regions of the parameter space.

The author thanks the Inter University Centre for Astronomy and Astrophysics, Pune, India for hospitality while this work was completed. This work is supported by the J. C. National Bose Fellowship of the Department of Science and Technology, and by the Council of Scientific and Industrial Research, India under Project No. (03)(1220)/12/EMR-II.

[1] R. Beckmann, F. Karsch and D. E. Miller, Phys. Rev. Lett. **43**, 1277 (1979) [Erratum-ibid. **45**, 1306 (1980)]; Phys. Rev. **A25**, 561 (1982); F. Karsch and D. E. Miller, *ibid***22**, 1210 (1980).
[2] C. Aragao De Carvalho and S. Goulart Rosa, J. Phys. A **13**, 3233 (1980); C. A. Aragao De Carvalho and S. Goulart Rosa, J. Phys. A **13**, 989 (1980); H. O. da Frota and S. Goulart Rosa, Jr., J. Phys. A **15**, 2221 (1982).
[3] H. E. Haber and H. A. Weldon, Phys. Rev. Lett. **46**, 1497 (1981); H. E. Haber and H. A. Weldon, Phys. Rev. D **25**, 502 (1982).
[4] S. Singh and P. N. Pandita, Phys. Lett. A **92**, 65 (1982); Phys. Rev. A **28**, 1752 (1983);
[5] K. Huang, *Statistical Mechanics*, (Wiley, NY, 1987), 2nd ed., pp. 154-7.
[6] P. T. Landsberg and J. Dunning-Davies, Phys. Rev. **138**, A1049 (1965).
[7] M. Grether, M. de Llano and G. A. . Baker, Phys. Rev. Lett. **99**, 200406 (2007) [arXiv:0706.2833 [cond-mat.supr-con]].
[8] M. M. Nieto, J. Math. Phys. **11**, 1346 (1970).
[9] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions*, (Dover, New York, 1972), p. 374ff.
[10] J. D. Gunton and M. J. Buckingham, Phys. Rev. **166**, 152 (1968).